

Is endoscopist-directed sedation for endoscopic retrograde cholangiopancreatography with balanced propofol solution (BPS) practical and safe? A prospective study from a tertiary care centre in India

Hameed Raina¹, Anukalp Prakash¹, Umaymah Ashraf¹, Manzoor Raina²

¹Department of Gastroenterology and Hepatology, Paras Hospital, Gurugram, India

²Department of Clinical Biochemistry, SKIMS, Soura, India

Gastroenterology Rev 2021; 16 (4): 324–329
DOI: <https://doi.org/10.5114/pg.2021.111764>

Key words: sedation, endoscopic retrograde cholangiopancreatography, balanced propofol solution.

Address for correspondence: Hameed Raina MBBS, MD, DM (PhD), Department of Gastroenterology and Hepatology, Paras Hospital, Gurugram, India, e-mail: hameedraina49@gmail.com

Abstract

Introduction: Endoscopist-directed, nurse-administered sedation using propofol, midazolam, and fentanyl for endoscopic retrograde cholangiopancreatography (ERCP) is being utilized worldwide. However, this is not usually employed in India by endoscopists.

Aim: To assess the efficacy, acceptability, and safety of this sedation in low to moderate risk patients undergoing ERCP.

Material and methods: This was a prospective study involving 500 patients with any indication for ERCP. The sedation was given by trained nurses.

Results: The sedative dosages per patient were as follows: propofol = 90 ±20 mg, fentanyl 0.75 ±0.25 mg (range: 0.25–1.00 mg), and midazolam 2 ±0.5 mg (range: 1–3 mg). Ninety-seven percent of patients achieved Richmond agitation sedation score of ≥ –3, and 96.8% achieved Gloucester comfort score of ≤ 2. 4.22% of the patients had mild adverse events (AE), and 2.11% had moderate AE. Two (0.4%) patients required intubation and intensive care unit admission. Mean recovery time was 15.3 min. 98.3% of the endoscopists were satisfied with the sedation achieved. 31.2% of the patients remembered the procedure. 93% of the patients were satisfied with the type of sedation. 92.75% of ERCPs were successful. 7.59% had procedure-related complications: in the form PEP – 5.290%, significant bleeding – 1.08%, and perforation – 0.43%.

Conclusions: Endoscopist-directed, nurse-administered sedation for ERCP with balanced propofol solution is practical, efficient, time saving, safe, and acceptable to patients.

Introduction

Endoscopist-directed nurse-administered sedation (EDNAS) using propofol and fentanyl in patients undergoing endoscopic retrograde cholangiopancreatography (ERCP) is being increasingly utilized worldwide. However, this method of sedation is not usually employed in India by endoscopists due to concerns surrounding its efficacy and safety. Propofol was introduced in the 1980s. It has become well-known for moderate sedation use for GI endoscopic procedures and its use has been increasingly adopted during the past decade [1]. It provides additional benefits including reduced time to induction of sedation, shorter duration of action, faster

patient recovery time, higher post-procedure patient satisfaction, and greater patient willingness to repeat endoscopy in the future [2]. Propofol-based sedation has a similar rate of adverse events to other forms of sedation. It can be safely administered with fixed doses of benzodiazepines and/or opioids to enhance hypnotic and sedative effects [3]. The combination of propofol with other sedative agents leads to a reduction in the propofol dose, thereby improving its safety profile [4]. This moderate sedation regimen is known as “balanced propofol solution” (BPS). In BPS, a benzodiazepine, propofol, and an opioid are administered in a single dose. This is then complemented by small incremental doses of propofol (10–20 mg/push) to obtain a target

level of moderate sedation [5]. However, higher doses of propofol can induce deep sedation leading to respiratory depression and apnoea [6]. Furthermore, there is currently no known propofol antagonist. Moreover, propofol product labelling states that “propofol should be administered only by persons trained in the administration of general anaesthesia” [7]. Unfortunately, there are few anaesthetists in developing countries, particularly in India.

No major study regarding EDNAS using propofol with midazolam/fentanyl for ERCP from India is available.

Aim

Hence, the aim of this study was to assess the efficacy, acceptability, and safety of this type of sedation in low to moderate risk patients undergoing ERCP in a developing nation like India where resources are limited for general anaesthesia.

Material and methods

This study was conducted at a single tertiary hospital in Paras Hospital, Gurugram, India. It was a prospective study involving 500 patients with an ASA score of I–III, presenting with any indication for ERCP. Informed consent was taken from all patients. A total of 500 patients (348 male and 152 females) with a mean age of 57.3 years (range: 20–87) were recruited from August 2016 to July 2019. The following patients were excluded:

1. Those who did not consent for the study.
2. Those in ASA class IV and V.
3. Those with difficult airways, e.g. with high Mallampati score, short neck, obesity, etc.
4. Those with a history of failed ERCP or non-cooperation or requiring double stenting.
5. Psychiatric or irritable patients.
6. Those below 18 or above 70 years of age (Figure 1).

The sedation was given by the trained nurses posted in the endoscopy OT. Intra-procedural and post-procedure (AEs) adverse events due to sedation were recorded by the doctors. The endoscopist was also responsible for directing the provision and dosing of the BPS. Complete cardiopulmonary resuscitation equipment and medications were available within the endoscopy unit. Endoscopy doctors and nurses were necessarily basic and advanced cardiac life-support certified.

In this study, we evaluated patient-level demographic variables including age, gender, ASA classification, and indication for ERCP. We also evaluated clinically relevant patient outcome variables including the following: BPS drug dosages, pre-procedure and post-procedure oxygen saturation levels, pulse, sys-

tolic/diastolic blood pressures, and the need for mask bag ventilation or endotracheal intubation at any time during or immediately following ERCP. Aborted ERCP procedures due to a sedation-related adverse event, any ICU admission, and mortality were also recorded. Institutional Ethics Committee approval was obtained for this study.

Statistical analysis

Descriptive statistics for all parameters were calculated: For continuous variables this included mean, standard deviation (SD), median, minimum, maximum, and range values. For non-continuous variables, counts and percentages were reported. A previously determined value of $p < 0.05$ was considered statistically significant.

Results

From 1 August 2016 to 30 October 2018, 500 patients undergoing ERCP received endoscopist-directed sedation.

Among them 150 (30%) were alcoholics, 225 (45%) were smokers, 348 (69.6%) were males, 152 (30.4%) were females (Table I), 115 (23%) were American Society of Anaesthesiologists (ASA) I, 320 (64%) were ASA II and 65 (13%) were ASA III with PS 1–2. The mean age was 57.3 years (Figure 2).

Indications for ERCP by the endoscopist-directed, nurse-administered (EDNA) sedation included 415 (83%) choledocholithiasis, 45 (9%) benign CBD strictures, 30 (6%) malignant obstructive jaundice (OJ), and 10 (2%) bile duct injuries (Table II).

Sedative dosages (mean \pm standard deviation) per patient were propofol = 90 ± 20 mg, fentanyl 0.75 ± 0.25 mg (range: 0.25–1.00 mg), and midazolam 2 ± 0.5 mg (range: 1–3 mg). Mean injection to sedation time was 2 min (range: 1–4 min). 97% of patients achieved a Richmond agitation sedation score (RASS) of ≥ -3 (Figure 3) and 96.8% achieved a Gloucester comfort score (GCS) of ≤ 2 (Figure 4) during the procedure. 4.22% of the patients had mild adverse events and 2.11% had moderate AE in the form of nausea, hypotension, breathing difficulty, or hypoxemia. Two (0.4%) patients required intubation and ICU admission. The remaining patients resolved all after conservative management in the recovery room itself. There were 3 (0.6%) aborted ERCPs due to moderate sedation-related AE. There was no mortality secondary to sedation-related AE (Figure 5). The mean recovery time was 15.3 min. 98.3% of the endoscopists were satisfied with the sedation achieved during procedure. 31.2% of the patients remembered the procedure. 93% of the patients were satisfied with the type of sedation, and 86.8% were ready for a re-




Figure 1. Flow chart

Table I. Demographics

Parameter	N	%
ASA I	115	23
ASA II	320	64
ASA III	65	13
Males	348	69.6
Females	152	30.4
Alcoholics	150	30
Smokers	225	45
Choledocholithiasis	415	83
Benign CBD stricture	45	9
Malignant OJ	30	6
Bile duct injury	10	2




Figure 2. American Society of Anaesthesiologists (ASA) classification of patients

Table II. Results of endoscopist direct nurse administered sedation with balanced propofol solution in ERCP

Drug	Result Mean dose ± mean deviation or %
Propofol	90 ±20 mg
Fentanyl	0.75 ±0.25 mg
Midazolam	2 ±0.5 mg
RAAS ≥ -3	97%
GCS ≤ 2	96.8%
Mild AE	4.22%
Moderate AE	2.11%
Severe AE	0.4%
Aborted procedures due to sedation-related AE	0.6%
Mean recovery time	15.3 min
Aldrete score ≥ 9/10	93%
% of patients satisfied with sedation	93%
% of patients ready for repeat procedure with same sedation	86.8%
% of endoscopists satisfied with sedation	98.3%
% of patients who remembered the procedure	31.2%




Figure 5. Adverse events

peat procedure, if indicated, under the same sedation (Figure 6).

The outcomes of ERCPs: Among 497 ERCPs done in patients under EDNAS, 461 (92.75%) were successful and 36 (7.24%) were unsuccessful due to techni-




Figure 3. Richmond Agitation Sedation Score (RASS)




Figure 4. Gloucester Comfort Score (GCS)




Figure 6. Patients' and endoscopists' experience with EDNAS

cal reasons other than sedation related. 35 (7.59%) had procedure-related complications: in the form of post-ERCP pancreatitis (PEP) in 24 (5.290%); significant bleeding requiring treatment in 5 (1.08%); perforation in 2 (0.43%); and others like ascending cholangitis in

4 (0.86%) patients. However, none had procedure-related mortality.

Discussion

There is ongoing controversy and lack of consensus regarding the safety and efficacy of endoscopist-directed BPS in the absence of monitored anaesthesia care during GI endoscopy procedures because it has a narrow therapeutic index and can cause irreversible respiratory depression.

No major study regarding EDNAS using propofol with midazolam/fentanyl for ERCP from India is available. Our study is one of the first such studies. We selected about 500 patients with ASA score of I to III. The majority of our patients had moderate risk (ASA II) due to comorbidities/jaundice.

The mean dosage of propofol was 90 ± 20 mg, fentanyl 0.75 ± 0.25 mg, and midazolam 2 ± 0.5 mg, which is less than used by Nathan *et al.* [8]. However, Clarke *et al.* evaluated 15,268 colonoscopies in which the mean propofol dose used was 60 mg, administered in combination with midazolam and fentanyl [9].

Our sedation protocol showed high efficacy; 97% of our patients achieved Richmond agitation sedation score (RASS) of ≥ -3 .

Mean recovery time was 15.3 min. This is lower than achieved by Sathanathan *et al.* in their study (23 min) [10]. Rapid recovery and return to baseline behaviour allowed us to quickly discharge patients or move them to the ward from the endoscopy unit after the procedures. This saved a lot of time and increased the number of procedures done per day.

4.22% and 2.11% of the patients had mild and moderate AE, respectively, in the form of hypotension, breathing difficulty, bradycardia, or hypoxaemia. 0.4% had severe AE and required intubation and ICU admission. 0.6% had aborted ERCPS due to moderate sedation-related AE. These percentages of AE are much lower than in previous studies. Quaid *et al.* reported 2–16% of sedation related AE [11]. Rex *et al.* reported in their study that 0.1% of their patients needed bag and mask ventilation, 11 patients required intubation, and 4 patients died [12].

98.3% of the endoscopists were satisfied with the sedation achieved. This is the same as seen in other studies. Physician satisfaction with propofol sedation was about 95% in 2 colonoscopy studies as reported by Ulmer *et al.* and Hansen *et al.* [13, 14].

The mean recovery time was 15.3 min. This is lower than that achieved by Sathanathan *et al.* in their study (23 min) [10].

31.2% of the patients had some memory of the procedures done to them, which is less than with midazol-

am or midazolam/fentanyl combination. Two colonoscopy trials found that there was 13% more memory of the procedures sedated with midazolam/narcotics than with propofol and 10% fewer were satisfied with their sedation [11].

93% of the patients were satisfied with the type of sedation, and 86.8% were ready for a repeat procedure, if indicated, under the same the sedation. The same results were shown by other studies.

Endoscopist-directed BPS may have substantial economic advantages; the financial burden of anaesthesiologist services for GI endoscopic procedures is estimated to be \$5 billion annually in the USA [15]. Furthermore, a lot of time is saved using EDNAS. This will increase the number of endoscopic procedures that can be done in the hospital. This is very important in a country like ours where the demand always exceeds the availability.

The outcomes of ERCP under EDNAS are similar to those done under anaesthetist-directed sedation with lower or similar complication rates [16]. Hence, our study proves that endoscopist-directed sedation does not affect the efficiency of the doctor.

Conclusions

Endoscopist-directed nurse-administered sedation for ERCP is easy, time saving, safe, and acceptable to patients when the endoscopists and nursing staff are adequately trained and strict patient selection criteria are used.

Acknowledgments

We wish to thank our endoscopy nursing staff for their support and active participation in this study. We owe this study to all the patients who participated, without whom this concept would never have materialized.

Conflict of interest

The authors declare no conflict of interest.

References

- Lewis JR, Cohen LB. Update on colonoscopy preparation, premedication and sedation. *Exp Rev Gastroenterol Hepatol* 2013; 7: 77-87.
- Shafer A, Doze VA, Shafer SL, White PF. Pharmacokinetics and pharmacodynamics of propofol infusions during general anesthesia. *Anesthesiology* 1988; 69: 348-56.
- Roseveare C, Seavell C, Patel P, et al. Patient-controlled sedation and analgesia, using propofol and alfentanil, during colonoscopy: a prospective randomized controlled trial. *Endoscopy* 1998; 30: 768-73.
- Cohen LB, Hightower CD, Wood DA, et al. Moderate level sedation during endoscopy: a prospective study using low-dose

- propofol, meperidine/fentanyl, and midazolam. *Gastrointest Endosc* 2004; 59: 795-803.
5. Vargo JJ, Cohen LB, Rex DK, Kwo PY. American Association for the Study of Liver Diseases, American College of Gastroenterology, American Gastroenterological Association, American Society for Gastrointestinal Endoscopy. Position statement. Nonanesthesiologist administration of propofol for GI endoscopy. *Gastroenterology* 2009; 137: 2161-7.
 6. Miner JR, Burton JH. Clinical practice advisory: emergency department procedural sedation with propofol. *Ann Emerg Med* 2007; 50: 182-7.
 7. AstraZeneca. Novaplus Diprivan 1% Propofol Injectable Emulsion Product-manufactures Recommendations. [cited 2009 Nov 06]. Available from: <http://www1.astrazeneca-us.com/pi/diprivan.pdf>.
 8. Nathan JH, Klein A, Gralnek IM, Khamaysi I. Endoscopist-directed balanced propofol sedation is safe and effective in patients undergoing outpatient colonoscopy. *J Dig Endosc* 2015; 6: 158-62.
 9. Clarke AC, Chiragakis L, Hillman LC, Kaye GL. Sedation for endoscopy: the safe use of propofol by general practitioner sedationists. *Med J Australia* 2002; 176: 158-61.
 10. Sathananthan D, Young E, Nind G, et al. Assessing the safety of physician-directed nurse-administered propofol sedation in low-risk patients undergoing endoscopy and colonoscopy. *Endosc Int Open* 2017; 5: E110-5.
 11. Quaid KR Mc, Laine L. A systematic review and meta-analysis of randomized, controlled trials of moderate sedation for routine endoscopic procedures. *Gastrointest Endosc* 2008; 67: 910-23.
 12. Rex DK, Deenadayalu VP, Eid E, et al. Endoscopist-directed administration of propofol: a worldwide safety experience. *Gastroenterology* 2009; 137: 1229-37.
 13. Ulmer BJ, Hansen JJ, Overley CA. Propofol versus midazolam/fentanyl for outpatient colonoscopy: administration by nurses supervised by endoscopists. *Clin Gastroenterol Hepatol* 2003; 1: 425-32.
 14. Hansen JJ, Ulmer BJ, Rex DK. Technical performance of colonoscopy in patients sedated with nurse-administered propofol. *Am J Gastroenterol* 2004; 99: 52-56.
 15. Aisenberg J, Cohen LB, Piorkowski JD Jr. Propofol use under the direction of trained gastroenterologists: an analysis of the medicolegal implications. *Am J Gastroenterol* 2007; 102: 707-13.
 16. Hormati A, Aminnejad R, Saeidi M, et al. Prevalence of anesthetic and gastrointestinal complications of endoscopic retrograde cholangiopancreatography. *Anesth Pain Med* 2019; 9: e95796.

Received: 7.12.2020

Accepted: 30.01.2021